



Astrocyte Medium – Glucose Free (AM-GF)

Catalog #1801-GF

Product Description

Astrocyte Medium – glucose free (AM-GF), when used with Astrocyte Growth Supplement (AGS), FBS and additional glucose (Cat #0893) is a medium designed for optimal growth of normal astrocytes *in vitro*. It is a sterile, liquid medium which contains essential and non-essential amino acids, vitamins, organic and inorganic compounds, hormones, growth factors, trace minerals and a low concentration of fetal bovine serum (2%). The medium is HEPES and bicarbonate buffered and has a pH of 7.4 when equilibrated in an incubator with an atmosphere of 5% CO₂/95% air. The medium is formulated (quantitatively and qualitatively) to provide an optimally balanced nutritional environment that selectively supports the growth of normal astrocytes *in vitro*.

AM-GF does not contain glucose, an important source of energy for cell growth. AM-GF should be supplemented with Glucose Solution (GS; Cat #0893) prior to use. The recommended glucose concentration is 1g/L but may vary with experimental conditions.

Components

AM-GF consists of 500 mL of basal medium, 10 mL of FBS (Cat. #0010), 5 mL of Astrocyte Growth Supplement (AGS, Cat. #1852) and 5 mL of penicillin/streptomycin solution (P/S, Cat. #0503). *Note: FBS, AGS and P/S are not pre-mixed in AM-GF; they must be added separately to make the complete AM-GF.*

NOTE: AM-GF should be supplemented with Glucose Solution (GS; Cat #0893) prior to use. Glucose Solution is not included and should be purchased separately.

Product Use

AM-GF is for research use only. It is not approved for human or animal use, or for application in *in vitro* diagnostic procedures.

Storage

Store the basal medium at 4°C and the AGS, FBS and P/S solution at -20°C. Protect from light.

Shipping

Basal medium: room temperature. Supplements: dry ice.

Instructions for Use

Thaw AGS, FBS and P/S solution at 37°C. Gently tilt the tubes several times to ensure the contents are completely mixed before adding to the medium. Spray the medium bottle and tubes with 70% ethanol, and wipe to remove excess liquid. In a sterile field, remove the caps without touching the interior threads with fingers. Add AGS, FBS and P/S solution to the medium and mix well. Since several components are light-labile, the medium should not be exposed to light

Rev. 0

for extended periods. We do not recommend warming medium in a 37°C water bath prior to use. When stored in the dark at 4°C, the reconstituted medium is stable for one month.

Caution: If handled improperly, some components of the medium may present a health hazard. Take appropriate precautions when handling it, including the wearing of protective clothing and eyewear. Dispose of properly.