

3D-Preadipocyte Spheroid Differentiation Medium (3D-PADM)

Catalog #3D-7221

Product Description

3D-Preadipocyte Spheroid Differentiation Medium (3D-PADM) is a specialized medium designed for the differentiation of human preadipocyte spheroids *in vitro* in conjunction with 3D-preadipocyte spheroid differentiation supplement (3D-PAdDS, Cat. #3D-7232). 3D-PADM is a sterile, viscous medium which contains essential and non-essential amino acids, vitamins, organic and inorganic compounds, and trace minerals. The medium is bicarbonate buffered and has a pH of 7.4 when equilibrated in an incubator with an atmosphere of 5% CO₂/95% air.

Components

3D-PADM consists of 200 mL of basal medium, 2 mL of 3D-preadipocyte spheroid differentiation supplement (3D-PAdDS, Cat. #3D-7232), 10 mL of fetal bovine serum (FBS, Cat. #0010) and 2 mL penicillin/streptomycin solution (P/S, Cat. #0583). *Note: 3D-PAdDS, FBS and P/S are not pre-mixed in the 3D-PADM; they must be added separately to make the complete medium.*

Product Use

3D-PADM is for research use only. It is not approved for human or animal use, or for application in *in vitro* diagnostic procedures.

Storage

Store the basal medium at 4°C and the 3D-PAdDS, FBS and P/S solution at -20°C. Protect from light.

Shipping

Basal medium: room temperature. Supplements: dry ice.

Instructions for Use

Thaw 3D-PAdDS, FBS and P/S solution at 37°C. Gently tilt the tubes several times to ensure the contents are completely mixed before adding to the basal medium. Spray the medium bottle and tubes with 70% ethanol and wipe to remove excess liquid. In a sterile field, remove the caps without touching the interior threads with fingers. Add 3D-PAdDS, FBS and P/S solution to the medium and mix carefully. Gentle mixing of the viscous 3D culture medium is recommended to avoid bubble formation. Since several components are light-labile, the medium should not be exposed to light for extended periods. We do not recommend warming medium in a 37°C water bath prior to use. When stored in the dark at 4°C, the reconstituted medium is stable for one month.

Caution: If handled improperly, some components of the medium may present a health hazard. Take appropriate precautions when handling it, including the wearing of protective clothing and eyewear. Dispose of properly.