



Cardiac Myocyte Medium – Glucose Free (CMM-GF)

Catalog #6201-GF

Product Description

Cardiac Myocyte Medium – glucose free (CMM-GF), when used with Cardiac Myocyte Growth Supplement (CMGS), FBS and additional glucose (Cat #0893) is a medium designed for optimal growth of normal cardiac myocytes *in vitro*. It is a sterile, liquid medium which contains essential and non-essential amino acids, vitamins, organic and inorganic compounds, hormones, growth factors, trace minerals and a low concentration of fetal bovine serum (5%). The medium is HEPES and bicarbonate buffered and has a pH of 7.4 when equilibrated in an incubator with an atmosphere of 5% CO₂/95% air. The medium is formulated (quantitatively and qualitatively) to provide an optimally balanced nutritional environment that selectively supports the growth of cardiac myocytes *in vitro*.

CMM-GF does not contain glucose, an important source of energy for cell growth. CMM-GF should be supplemented with Glucose Solution (GS; Cat #0893) prior to use. The recommended glucose concentration is 1g/L but may vary with experimental conditions.

Components

CMM-GF consists of 500 ml of basal medium, 25 mL of fetal bovine serum (FBS, Cat. #0025), 5 mL of Cardiac Myocyte Growth Supplement (CMGS, Cat. #6252) and 5 mL of penicillin/streptomycin solution (P/S, Cat. #0503). *Note: FBS, CMGS and P/S are not pre-mixed in CMM-GF; they must be added separately to make the complete CMM-GF.*

NOTE: CMM-GF should be supplemented with L-glutamine prior to use. L-glutamine is not provided with this kit and must be purchased separately.

Product Use

NOTE: CMM-GF should be supplemented with Glucose Solution (GS; Cat #0893) prior to use. Glucose Solution is not included and should be purchased separately.

Storage

Store the basal medium at 4°C and the CMGS, FBS and P/S solution at -20°C. Protect from light.

Shipping

Basal medium: room temperature. Supplements: dry ice.

Instructions for Use

Thaw CMGS, FBS and P/S solution at 37°C. Gently tilt the tubes several times to ensure the contents are completely mixed before adding to the medium. Spray the medium bottle and tubes with 70% ethanol, and wipe to remove excess liquid. In a sterile field, remove the caps without touching the interior threads with fingers. Add CMGS, FBS and P/S solution to the medium and mix well. Since several components are light-labile, the medium should not be exposed to light.

Rev. 0

for extended periods. We do not recommend warming medium in a 37°C water bath prior to use. When stored in the dark at 4°C, the reconstituted medium is stable for one month.

Caution: If handled improperly, some components of the medium may present a health hazard. Take appropriate precautions when handling it, including the wearing of protective clothing and eyewear. Dispose of properly.